

AD-A091 815

PACIFIC-SIERRA RESEARCH CORP SANTA MONICA CA  
IMPROVED NOISE SUPPRESSION FOR MULTIBAND IMAGERY.(U)


F/G 9/2

UNCLASSIFIED

AUG 80 P J READY, D A ANDERSON, B J GAITLEY F49620-78-C-0059  
PSR-1028 AFOSR-TB-80-1176

MR

END  
DATE  
FILMED  
1-8  
DTIC



MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

2  
AFOSR-TR. 80 - 1134

PSR Report 1028

LEVEL

8

IMPROVED NOISE SUPPRESSION FOR MULTIBAND IMAGERY

P. J. Ready  
D. A. Anderson  
B. J. Gaitley

August 1980

DTIC  
ELECTE  
NOV 13 1980  
C

Final Technical Report  
Contract F49620-78-C-0059

Sponsored by  
Air Force Office of Scientific Research  
Bolling Air Force Base  
Washington, D.C. 20332



PACIFIC-SIERRA RESEARCH CORP.

1486 Cloverfield Blvd. • Santa Monica, California 90404

BDC FILE COPY

80 11 06 097

Approved for public release;  
distribution unlimited.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER <b>AFOSR-TR- 80 - 1134</b>	2. GOVT ACCESSION NO. <b>AD-A092 815</b>	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) <b>IMPROVED NOISE SUPPRESSION FOR MULTIBAND IMAGERY</b>		5. TYPE OF REPORT & PERIOD COVERED <b>Final</b>
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) <b>P. J. Ready D. A. Anderson B. J. Gaitley</b>		8. CONTRACT OR GRANT NUMBER(s) <b>F49620-78-C-0059</b>
9. PERFORMING ORGANIZATION NAME AND ADDRESS <b>Pacific-Sierra Research corporation 1456 Cloverfield Blvd. Santa Monica, California 90404</b>		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS <b>61102F 2304/A6</b>
11. CONTROLLING OFFICE NAME AND ADDRESS <b>Air Force Office of Scientific Research/NM Bolling Air Force Base Washington DC 20332</b>		12. REPORT DATE <b>August 1980</b>
		13. NUMBER OF PAGES <b>25</b>
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) <b>UNCLASSIFIED</b>
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report)  <b>Approved for public release; Distribution unlimited</b>		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)  <b>Image processing SNR improvement Multiband images Karhunen-Loeve transformation</b> <i>This research has</i>		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) <b>Develop and test an approach to improvement of noise degraded multiband imagery based on a class dependent Karhunen-Loeve (K-L) transformation and transform domain spatial filtering. The technique is attractive because of its tendency to preserve the high spatial frequency information (e.g., edges in the picture as compared with conventional minimum mean square error linear filtering of the original bands. Multiband images (especially color and multispectral) characteristically exhibit significant class structure, and within a particular class show high spectral correlations. The filtering</b>		

DD FORM 1 JAN 73 1473 EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

~~UNCLASSIFIED~~

A technique developed during this research exploits both the class structure and the intraclass correlation to achieve noise suppression.

11

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

14

PSR 1028

8

IMPROVED NOISE SUPPRESSION FOR MULTIBAND IMAGERY

9 Final Report

10

P. J. Ready  
D. A. Anderson  
B. J. Gaitley

DTIC  
ELECTED  
NOV 13 1980

11

August 1980

12 27

16 2304

17 A6

18 AFOSR

Final Technical Report  
Contract F49620-78-C-0059

15

19 7K-80-1134

Sponsored by  
Air Force Office of Scientific Research  
Bolling Air Force Base  
Washington, D.C. 20332



PACIFIC-SIERRA RESEARCH CORP.

1456 Cloverfield Blvd. • Santa Monica, California 90404

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFSC)  
NOTICE OF TRANSMITTAL TO DDC  
This technical report has been reviewed and is  
approved for release IAW AFR 190-12 (7b).  
Distribution is unlimited.  
A. D. BLOSE  
Technical Information Officer

407411

R

## CONTENTS

### Section

1. INTRODUCTION ..... 1
2. DEFINITIONS ..... 1
3. DERIVATION OF THE MATRIX WIENER-HOPF EQUATION ..... 6
4. SOLUTION UNDER THE ASSUMPTION OF STATIONARITY AND  
NON-CAUSALITY ..... 16.
5. CLASS-DEPENDENT K-L TRANSFORMATION ..... 17

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A	

## FINAL TECHNICAL REPORT

### IMPROVED NOISE SUPPRESSION FOR MULTIBAND IMAGERY

#### 1.0 INTRODUCTION

This research has developed and tested an approach to improvement of noise degraded multiband imagery based on a class dependent Karhunen-Loeve (K-L) transformation and transform domain spatial filtering. The technique is attractive because of its tendency to preserve the high spatial frequency information (e.g., edges) in the picture as compared to conventional minimum mean square error linear filtering of the original bands. By multi-band we mean a set of images of the same scene as recorded in different optical wavelengths, at different times, or a combination of both. For example, an ordinary color photograph is composed of three bands--red, green, and blue. Multispectral imagery is another example for which the same scene is imaged in several bands ranging from the visible to the infrared. Color video (e.g., television) is an example of the combination of spectral and temporal bands.

Multi-band images (especially color and multispectral) characteristically exhibit significant class structure, and within a particular class shows high spectral correlations. The filtering technique developed during this research exploits both the class structure and the intra class correlation to achieve noise suppression.

#### 2.0 DEFINITIONS

The multi-band image is viewed as a vector random process  $\underline{P}(x, y)$  where

$$\underline{P}(x, y) \triangleq \begin{bmatrix} P_1(x, y) \\ P_2(x, y) \\ \vdots \\ P_N(x, y) \end{bmatrix} \quad (1)$$



and  $P_i(x, y)$  is the amplitude of the  $i$ th image having spatial variables  $x$  and  $y$ . We define the  $N \times N$  spectral covariance matrix as

$$\underline{C}(x, y) \triangleq E\{\underline{P}(x, y) \underline{P}^t(x, y)\} \quad (2)$$

where  $E$  is the expectation,  $t$  indicates transpose, and  $E\{\underline{P}(x, y)\} = \underline{0}$  is assumed. We also assume spatial stationarity so that

$$\underline{C}(x, y) = \underline{C} . \quad (3)$$

The general stationary covariance matrix for the process is

$$\underline{K}(\tau_x, \tau_y) \triangleq E\{\underline{P}(x, y) \underline{P}^t(x + \tau_x, y + \tau_y)\} , \quad (4)$$

which includes the spatial covariance. The  $(i, j)$  the element of  $\underline{K}(\tau_x, \tau_y)$  is

$$k_{i,j}(\tau_x, \tau_y) = E\{P_i(x, y) P_j(x + \tau_x, y + \tau_y)\} . \quad (5)$$

To a good approximation,

$$k_{i,j}(\tau_x, \tau_y) = c_{i,j} R(\tau_x, \tau_y) \quad (6)$$

where  $c_{i,j}$  is the  $(i, j)$ th element of  $\underline{C}$  and

$$R(\tau_x, \tau_y) \triangleq E\{P_i(x, y) P_i(x + \tau_x, y + \tau_y)\} / \sigma^2 \quad i = 1, 2, \dots, N \quad (7)$$

is the normalized scalar autocovariance function for all elements of  $\underline{P}(x, y)$ . That is, each band is assumed to have the same spatial autocovariance function  $R(\tau_x, \tau_y)$  and variance  $\sigma^2$ . This is verified experimentally as shown in Figure 1 where we show the (azimuthally averaged) autocovariance function for the red, green, and blue bands




Fig. 1--Azimuthal Average of Red, Green, and Blue Autocorrelation Functions.

of a "typical" test image.

From (4) and (6),

$$\underline{K}(\tau_x, \tau_y) = \underline{C} \underline{R}(\tau_x, \tau_y) . \quad (8)$$

With the above definitions and assumptions we next derive the general covariance matrix for the principle component vector  $\underline{\alpha}(x, y)$  ,

$$\underline{\alpha}(x, y) \triangleq \underline{T} \underline{P}(x, y) \quad (9)$$

where  $\underline{T}$  is the  $N \times N$  matrix of eigenvectors of  $\underline{C}$ . Thus,

$$\begin{aligned} \underline{K}_{\alpha}(\tau_x, \tau_y) &\triangleq E\{\underline{\alpha}(x, y) \underline{\alpha}^t(x + \tau_x, y + \tau_y)\} \\ &= E\{\underline{T} \underline{P}(x, y) \underline{P}^t(x + \tau_x, y + \tau_y) \underline{T}^t\} \\ &= \underline{T} \underline{K}(\tau_x, \tau_y) \underline{T}^t . \end{aligned} \quad (10)$$

But  $\underline{K}(\tau_x, \tau_y) = \underline{C} \underline{R}(\tau_x, \tau_y)$  . Thus (10) becomes

$$\begin{aligned} \underline{K}_{\alpha}(\tau_x, \tau_y) &= (\underline{T} \underline{C} \underline{T}^t) \underline{R}(\tau_x, \tau_y) \\ &= \underline{R}(\tau_x, \tau_y) \begin{bmatrix} \lambda_1 & & 0 \\ & \lambda_2 & \\ & & \ddots \\ 0 & & & \lambda_N \end{bmatrix} \\ &= \underline{R}(\tau_x, \tau_y) \underline{\lambda} \end{aligned} \quad (11)$$

since  $C$  is diagonalized by the orthogonal similarity transformation  $T C T^T$ . The  $\lambda_i$ ,  $i = 1, 2, \dots, N$  are the eigenvalues of  $C$  ordered such that  $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_N$ .

Equation (11) is a key result. It states that the principal component images are orthogonal, that their variances are distributed as the  $\lambda_i$ , and that their individual spatial autocovariance functions are

$$\begin{aligned} R_{i\alpha}(\tau_x, \tau_y) &\triangleq E\{\alpha_i(x, y) \alpha_i(x + \tau_x, y + \tau_y)\} \\ &= \lambda_i R(\tau_x, \tau_y) \quad i = 1, 2, \dots, N. \end{aligned} \quad (12)$$

That is, each  $R_{i\alpha}(\tau_x, \tau_y)$  is a scaled (by  $\lambda_i$ ) replica of the original spatial autocovariance function. The impact of this result is developed next.

### 3.0 DERIVATION OF THE MATRIX WIENER-HOPF EQUATION

To determine the filter generating the minimum variance linear estimate of a vector picture process received in the presence of additive white noise the following problem formulation is needed. Suppose  $\underline{P}(\underline{x})$  is the n-element vector picture process, with  $\underline{x} = (x, y)$  the generic point on the picture surface, and  $\underline{n}(\underline{x})$  is the additive white observation noise. Then, the problem is to find a matrix spatial impulse response  $\underline{H}^{(0)}(\underline{x}, \underline{y})$  such that

$$\begin{aligned} \min_{\text{all } \underline{H}(\underline{x}, \underline{y})} E\{ \|\underline{P}(\underline{x}) - \int_A \underline{H}(\underline{x}, \underline{y}) [\underline{P}(\underline{y}) + \underline{n}(\underline{y})] d\underline{y}\|^2 \} \\ = E\{ \|\underline{P}(\underline{x}) - \int_A \underline{H}^{(0)}(\underline{x}, \underline{y}) [\underline{P}(\underline{y}) + \underline{n}(\underline{y})] d\underline{y}\|^2 \} \end{aligned} \quad (13)$$

for all  $\underline{x}$  in the 2-dimensional area A defined by the picture. Here "all  $\underline{H}(\underline{x}, \underline{y})$ " means all square-integrable  $\underline{H}(\underline{x}, \underline{y})$ , i.e., all matrices  $\underline{H}(\underline{x}, \underline{y}) = [\{h_{ij}(\underline{x}, \underline{y})\}_{i,j=1}^n]$  such that

$$\begin{aligned} \int_A \int_A \|\underline{H}(\underline{x}, \underline{y})\|^2 d\underline{x} d\underline{y} \triangleq \int_A \int_A \sum_{i,j=1}^n h_{ij}^2(\underline{s}, \underline{y}) d\underline{x} d\underline{y} \\ < +\infty \end{aligned}$$

For convenience,  $\underline{P}(\underline{x}) + \underline{n}(\underline{x})$ , the received picture process, will be denoted by  $\underline{r}(\underline{x})$ :

$$\underline{r}(\underline{x}) \triangleq \underline{P}(\underline{x}) + \underline{n}(\underline{x}) .$$

Here

$$\begin{aligned} E[\underline{r}(\underline{x})\underline{r}^t(\underline{y})] &= E[\underline{p}(\underline{x})\underline{p}^t(\underline{y})] \\ + E[\underline{n}(\underline{x})\underline{n}^t(\underline{y})] &= E[\underline{p}(\underline{x})\underline{p}^t(\underline{y})] \\ + \frac{1}{2}N_0\delta(\underline{x}-\underline{y})\underline{I} & \end{aligned} \quad (14)$$

$\{\underline{I} = \{\delta_{ij}\}$  and for any  $\underline{v}$ ,  $\underline{v}^t$  is the transpose of  $\underline{v}\}$ .

Now it will be verified that if  $\underline{H}^{(0)}(\underline{x}, \underline{y})$  satisfies a matrix analogue of the scalar Wiener-Hopf integral equation  $\underline{H}^{(0)}(\underline{x}, \underline{y})$  does indeed provide the minimization of mean square error described by (1). It will be assumed that

$$E\{[\underline{p}(\underline{x}) - \int_A \underline{H}^{(0)}(\underline{x}, \underline{y})\underline{r}(\underline{y})d\underline{y}]\underline{r}^t(\underline{z})\} = 0, \text{ all } \underline{z} \text{ in } A \quad (15)$$

Note that (15) is equivalent to

$$\underline{R}_{\underline{p}, \underline{r}}(\underline{x}, \underline{z}) - \int_A \underline{H}^{(0)}(\underline{x}, \underline{y})\underline{R}_{\underline{r}, \underline{r}}(\underline{y}, \underline{z})d\underline{y} = 0 \quad (16)$$

where

$$\underline{R}_{\underline{p}, \underline{r}}(\underline{x}, \underline{z}) \triangleq E[\underline{p}(\underline{x})\underline{r}^t(\underline{z})]$$

and

$$\underline{R}_{\underline{r}, \underline{r}}(\underline{y}, \underline{z}) \triangleq E[\underline{r}(\underline{y})\underline{r}^t(\underline{z})].$$

Thus, (15) is, in the form (16), a matrix Wiener-Hopf equation.

Write the arbitrary square integrable  $\underline{H}(\underline{x}, \underline{y})$  as a perturbation of  $\underline{H}^{(0)}(\underline{x}, \underline{y})$ :

$$\underline{H}(\underline{x}, \underline{y}) = \underline{H}^{(0)}(\underline{x}, \underline{y}) = \underline{\Delta}(\underline{x}, \underline{y}) .$$

Then, the following sequence of equalities holds, where the operator "tr" is defined by the equation

$$\text{tr}(\{m_{ij}\}) \stackrel{\Delta}{=} \sum_{i=1}^n m_{ii}, \text{ all matrices } \{m_{ij}\}$$

$$\begin{aligned} & E\left\{\left\|\underline{P}(\underline{x}) - \int_A \underline{H}(\underline{x}, \underline{y}) \underline{r}(\underline{y}) d\underline{y}\right\|^2\right\} \\ &= E\left\{\text{tr}\left[\left(\underline{P}(\underline{x}) - \int_A \underline{H}^{(0)}(\underline{x}, \underline{y}) \underline{r}(\underline{y}) d\underline{y}\right)\left(\underline{P}(\underline{x}) - \int_A \underline{H}^{(0)}(\underline{x}, \underline{y}) \underline{r}(\underline{y}) d\underline{y}\right)^t\right]\right\} \\ &= E\left\{\text{tr}\left[\left(\underline{P}(\underline{x}) - \int_A \underline{H}^{(0)}(\underline{x}, \underline{y}) \underline{r}(\underline{y}) d\underline{y} - \int_A \underline{\Delta}(\underline{x}, \underline{y}) \underline{r}(\underline{y}) d\underline{y}\right)\right.\right. \\ &\quad \left.\left.\times \left(\underline{P}(\underline{x}) - \int_A \underline{H}^{(0)}(\underline{x}, \underline{y}) \underline{r}(\underline{y}) d\underline{y} - \int_A \underline{\Delta}(\underline{x}, \underline{y}) \underline{r}(\underline{y}) d\underline{y}\right)^t\right]\right\} \\ &= E\left\{\text{tr}\left[\left(\underline{P}(\underline{x}) - \int_A \underline{H}^{(0)}(\underline{x}, \underline{y}) \underline{r}(\underline{y}) d\underline{y}\right)\left(\underline{P}(\underline{x}) - \int_A \underline{H}^{(0)}(\underline{x}, \underline{y}) \underline{r}(\underline{y}) d\underline{y}\right)^t\right.\right. \\ &\quad \left.\left.- \left(\underline{P}(\underline{x}) - \int_A \underline{H}^{(0)}(\underline{x}, \underline{y}) \underline{r}(\underline{y}) d\underline{y}\right)\left(\int_A \underline{\Delta}(\underline{x}, \underline{y}) \underline{r}(\underline{y}) d\underline{y}\right)^t\right.\right. \\ &\quad \left.\left.- \left(\int_A \underline{\Delta}(\underline{x}, \underline{y}) \underline{r}(\underline{y}) d\underline{y}\right)\left(\underline{P}(\underline{x}) - \int_A \underline{H}^{(0)}(\underline{x}, \underline{y}) \underline{r}(\underline{y}) d\underline{y}\right)^t\right.\right. \\ &\quad \left.\left.- \left(\int_A \underline{\Delta}(\underline{x}, \underline{y}) \underline{r}(\underline{y}) d\underline{y}\right)\left(\int_A \underline{\Delta}(\underline{x}, \underline{y}) \underline{r}(\underline{y}) d\underline{y}\right)^t\right]\right\} \end{aligned}$$

$$\begin{aligned}
 & + E\{ \text{tr} [ - \left( \int_A \underline{\Delta}(\underline{x}, \underline{y}) \underline{r}(\underline{y}) d\underline{y} \right) (\underline{P}(\underline{x}) - \int_A \underline{H}^{(0)}(\underline{x}, \underline{y}) \underline{r}(\underline{y}) d\underline{y})^t \\
 & + \left( \int_A \underline{\Delta}(\underline{x}, \underline{y}) \underline{r}(\underline{y}) d\underline{y} \right) \left( \int_A \underline{\Delta}(\underline{x}, \underline{y}) \underline{r}(\underline{y}) d\underline{y} \right)^t ] \} \quad (17)
 \end{aligned}$$

Since the "tr" operator is linear and is invariant under matrix transposition, (17) becomes

$$\begin{aligned}
 & E\{ \| \underline{P}(\underline{x}) - \int_A \underline{H}(\underline{x}, \underline{y}) \underline{r}(\underline{y}) d\underline{y} \|^2 \} \\
 & = E\{ \text{tr} [ (\underline{P}(\underline{x}) - \int_A \underline{H}^{(0)}(\underline{x}, \underline{y}) \underline{r}(\underline{y}) d\underline{y}) (\underline{P}(\underline{x}) - \int_A \underline{H}^{(0)}(\underline{x}, \underline{y}) \underline{r}(\underline{y}) d\underline{y})^t ] \} \\
 & \quad - 2E\{ \text{tr} [ (\underline{P}(\underline{x}) - \int_A \underline{H}^{(0)}(\underline{x}, \underline{y}) \underline{r}(\underline{y}) d\underline{y}) \left( \int_A \underline{\Delta}(\underline{x}, \underline{y}) \underline{r}(\underline{y}) d\underline{y} \right)^t ] \} \\
 & \quad + E\{ \text{tr} [ \left( \int_A \underline{\Delta}(\underline{x}, \underline{y}) \underline{r}(\underline{y}) d\underline{y} \right) \left( \int_A \underline{\Delta}(\underline{x}, \underline{y}) \underline{r}(\underline{y}) d\underline{y} \right)^t ] \} \quad (18)
 \end{aligned}$$

But, first,

$$\begin{aligned}
 & E\{ \text{tr} [ (\underline{P}(\underline{x}) - \int_A \underline{H}^{(0)}(\underline{x}, \underline{y}) \underline{r}(\underline{y}) d\underline{y}) (\underline{P}(\underline{x}) - \int_A \underline{H}^{(0)}(\underline{x}, \underline{y}) \underline{r}(\underline{y}) d\underline{y})^t ] \} \\
 & = E\{ \| \underline{P}(\underline{x}) - \int_A \underline{H}^{(0)}(\underline{x}, \underline{y}) \underline{r}(\underline{y}) d\underline{y} \|^2 \} \quad (19)
 \end{aligned}$$



Moreover,

$$\begin{aligned}
 & E\{\text{tr}[(\underline{P}(\underline{x}) - \int_A \underline{H}^{(0)}(\underline{x}, \underline{y}) \underline{r}(\underline{y}) d\underline{y}) (\int_A \underline{\Delta}(\underline{x}, \underline{y}) \underline{r}(\underline{y}) d\underline{y})^t]\} \\
 &= E\{\text{tr}[\int_A [(\underline{P}(\underline{x}) - \int_A \underline{H}^{(0)}(\underline{x}, \underline{y}) \underline{r}(\underline{y}) d\underline{y}) \underline{r}^t(\underline{z})] \underline{\Delta}^t(\underline{x}, \underline{z}) d\underline{z}]\} \\
 &= \text{tr}\{\int_A E[(\underline{P}(\underline{x}) - \int_A \underline{H}^{(0)}(\underline{x}, \underline{y}) \underline{r}(\underline{y}) d\underline{y}) \underline{r}^t(\underline{z})] \underline{\Delta}^t(\underline{x}, \underline{z}) d\underline{z}\} \quad (20)
 \end{aligned}$$

Finally, using (14),

$$\begin{aligned}
 & E\{\text{tr}[(\int_A \underline{\Delta}(\underline{x}, \underline{y}) \underline{r}(\underline{y}) d\underline{y}) (\int_A \underline{\Delta}(\underline{x}, \underline{y}) \underline{r}(\underline{y}) d\underline{y})^t]\} \\
 &= E\{\text{tr}[(\int_A \underline{\Delta}(\underline{x}, \underline{y}) \underline{r}(\underline{y}) d\underline{y}) (\int_A \underline{r}^t(\underline{y}) \underline{\Delta}^t(\underline{x}, \underline{y}) d\underline{y})]\} \\
 &= E\{\text{tr}[\int_A \int_A \underline{\Delta}(\underline{x}, \underline{y}) \underline{r}(\underline{y}) \underline{r}^t(\underline{z}) \underline{\Delta}^t(\underline{x}, \underline{z}) d\underline{y} d\underline{z}]\} \\
 &= \text{tr}[\int_A \int_A \underline{\Delta}(\underline{x}, \underline{y}) E[\underline{r}(\underline{y}) \underline{r}^t(\underline{z})] \underline{\Delta}^t(\underline{x}, \underline{z}) d\underline{y} d\underline{z}] \\
 &= \text{tr}[\int_A \int_A \underline{\Delta}(\underline{x}, \underline{y}) E[\underline{P}(\underline{y}) \underline{P}^t(\underline{z})] \underline{\Delta}^t(\underline{x}, \underline{z}) d\underline{y} d\underline{z}]
 \end{aligned}$$

$$\begin{aligned}
 & + \operatorname{tr} \left[ \int_A \int_A \underline{\Delta}(\underline{x}, \underline{y}) \left( \frac{N_0}{2} \underline{I} \right) \underline{\Delta}^t(\underline{x}, \underline{z}) d\underline{y} d\underline{z} \right] \\
 & = E \left\{ \operatorname{tr} \left[ \left( \int_A \underline{\Delta}(\underline{x}, \underline{y}) \underline{P}(\underline{y}) d\underline{y} \right) \left( \int_A \underline{\Delta}(\underline{x}, \underline{y}) \underline{P}(\underline{y}) d\underline{y} \right)^t \right] \right\} \\
 & \quad + \operatorname{tr} \left[ (\operatorname{Area}(A)) \int_A \underline{\Delta}(\underline{x}, \underline{y}) \underline{\Delta}^t(\underline{x}, \underline{y}) d\underline{y} \right] \\
 & = E \left\{ \left\| \int_A \underline{\Delta}(\underline{x}, \underline{y}) \underline{P}(\underline{y}) d\underline{y} \right\|^2 \right\} + \operatorname{Area}(A) \int_A \left\| \underline{\Delta}(\underline{x}, \underline{y}) \right\|^2 d\underline{y} \quad (21)
 \end{aligned}$$

where

$$\left\| \underline{\Delta}(\underline{x}, \underline{y}) \right\|^2 = \sum_{i,j=1}^3 \Delta_{ij}^2(\underline{x}, \underline{y}) .$$

Hence, combining (18), (19), (20) and (21), there results

$$\begin{aligned}
 E \left\{ \left\| \underline{P}(\underline{x}) - \int_A \underline{H}(\underline{x}, \underline{y}) \underline{r}(\underline{y}) d\underline{y} \right\|^2 \right\} & = E \left\{ \left\| \underline{P}(\underline{x}) - \int_A \underline{H}^{(0)}(\underline{x}, \underline{y}) \underline{r}(\underline{y}) d\underline{y} \right\|^2 \right\} \\
 & + \operatorname{tr} \left\{ \int_A E \left[ \left( \underline{P}(\underline{x}) - \int_A \underline{H}^{(0)}(\underline{x}, \underline{y}) \underline{r}(\underline{y}) d\underline{y} \right) \underline{r}^t(\underline{z}) \right] \underline{\Delta}^t(\underline{x}, \underline{z}) d\underline{z} \right\} \\
 & + E \left\{ \left\| \int_A \underline{\Delta}(\underline{x}, \underline{y}) \underline{P}(\underline{y}) d\underline{y} \right\|^2 \right\} + \operatorname{Area}(A) \int_A \left\| \underline{\Delta}(\underline{x}, \underline{y}) \right\|^2 d\underline{y} \quad (22)
 \end{aligned}$$

Then, invoking the Wiener-Hopf equation (15), there results

$$\begin{aligned} E\{ \|\underline{P}(\underline{x}) - \int_A \underline{H}(\underline{x}, \underline{y}) \underline{r}(\underline{y}) d\underline{y}\|^2 \} &= E\{ \|\underline{P}(\underline{x}) - \int_A \underline{H}^{(0)}(\underline{x}, \underline{y}) \underline{r}(\underline{y}) d\underline{y}\|^2 \} \\ &+ E\{ \|\int_A \underline{\Delta}(\underline{x}, \underline{y}) \underline{P}(\underline{y}) d\underline{y}\|^2 \} + \text{Area}(A) \int_A \|\underline{\Delta}(\underline{x}, \underline{y})\|^2 d\underline{y} \end{aligned} \quad (23)$$

Thus, whenever for some  $i_0$  and  $j_0$

$$\int_A \Delta_{i_0 j_0}^2(\underline{x}, \underline{y}) d\underline{y} > 0,$$

it follows from (22) that

$$E\{ \|\underline{P}(\underline{x}) - \int_A \underline{H}(\underline{x}, \underline{y}) \underline{r}(\underline{y}) d\underline{y}\|^2 \} > E\{ \|\underline{P}(\underline{x}) - \int_A \underline{H}^{(0)}(\underline{x}, \underline{y}) \underline{r}(\underline{y}) d\underline{y}\|^2 \}$$

But, when for all  $i$  and  $j$

$$\int_A \Delta_{ij}^2(\underline{x}, \underline{y}) d\underline{y} = 0,$$

i.e., when

$$\int_A \|\underline{\Delta}(\underline{x}, \underline{y})\|^2 d\underline{y} = 0,$$

$$E\{ \|\underline{P}(\underline{x}) - \int_A \underline{H}(\underline{x}, \underline{y}) \underline{r}(\underline{y}) d\underline{y}\|^2 \} = E\{ \|\underline{P}(\underline{x}) - \int_A \underline{H}^{(0)}(\underline{x}, \underline{y}) \underline{r}(\underline{y}) d\underline{y}\|^2 \}$$

by (22),

Hence, not only does  $\underline{H}^{(0)}(\underline{x}, \underline{y})$  minimize  $E\{\|\underline{P}(\underline{x}) - \int_{\Lambda} \underline{H}(\underline{x}, \underline{y}) \underline{r}(\underline{y}) d\underline{y}\|^2\}$

but any minimizing  $\underline{H}(\underline{x}, \underline{y})$  has the property that

$$\begin{aligned} & \int_{\Lambda} \int_{\Lambda} \|\underline{H}(\underline{x}, \underline{y}) - \underline{H}^{(0)}(\underline{x}, \underline{y})\|^2 d\underline{x} d\underline{y} \\ &= \int_{\Lambda} \int_{\Lambda} \|\underline{\Delta}(\underline{x}, \underline{y})\|^2 d\underline{x} d\underline{y} \\ &= 0, \end{aligned}$$

i.e.,  $\underline{H}(\underline{x}, \underline{y})$  differs from  $\underline{H}^{(0)}(\underline{x}, \underline{y})$  by a null function.

Thus a solution of the Wiener-Hopf equation (15) (or (16)) not only minimizes the mean square error but it is unique in that respect to within a null function. This means, in addition, that if the Wiener-Hopf equation has one square-integrable solution it is unique to within a null function.

To see that the existence of a solution of the Wiener-Hopf equation is a necessary, as well as sufficient, condition for the existence of a minimizing choice of  $\underline{H}(\underline{x}, \underline{y})$  one must simply analyze equation (22). This equation is true independent of whether or not  $\underline{H}^{(0)}(\underline{x}, \underline{y})$  is a solution of the Wiener-Hopf equation. But if  $\underline{H}^{(0)}(\underline{x}, \underline{y})$  is not a solution of the Wiener-Hopf then set

$$\underline{\Delta}(\underline{x}, \underline{z}) = -\epsilon \underline{M}(\underline{x}, \underline{z}) \quad (24)$$

where  $\epsilon$  is an arbitrary positive number and

$$\underline{M}(\underline{x}, \underline{z}) \triangleq E[(\underline{P}(\underline{x}) - \int_{\Lambda} \underline{H}^{(0)}(\underline{x}, \underline{y}) d\underline{y}) \underline{r}^t(\underline{z})] \quad (25)$$

In this case,

$$\begin{aligned}
 & \text{tr} \left\{ \int_A E \left[ (\underline{P}(\underline{x}) - \int_A \underline{H}^{(0)}(\underline{x}, \underline{y}) \underline{r}(\underline{y}) d\underline{y}) \underline{r}^t(\underline{z}) \right] \underline{\Delta}^t(\underline{x}, \underline{y}) d\underline{z} \right\} \\
 &= -\epsilon \text{tr} \left\{ \int_A \underline{M}(\underline{x}, \underline{z}) \underline{M}^t(\underline{x}, \underline{z}) d\underline{z} \right\} \\
 &= -\epsilon \int_A \text{tr} \{ \underline{M}(\underline{x}, \underline{z}) \underline{M}^t(\underline{x}, \underline{z}) \} d\underline{z} \\
 &= -\epsilon \int_A \| \underline{M}(\underline{x}, \underline{z}) \|^2 d\underline{z} \tag{26}
 \end{aligned}$$

As a result of (26), (22) becomes

$$\begin{aligned}
 E \{ \| \underline{P}(\underline{x}) - \int_A \underline{H}(\underline{x}, \underline{y}) \underline{r}(\underline{y}) d\underline{y} \|^2 \} &= E \{ \| \underline{P}(\underline{x}) - \int_A \underline{H}^{(0)}(\underline{x}, \underline{y}) \underline{r}(\underline{y}) d\underline{y} \|^2 \} \\
 &- \epsilon \int_A \| \underline{M}(\underline{x}, \underline{z}) \|^2 d\underline{z} + E \{ \| \int_A (-\epsilon) \underline{M}(\underline{x}, \underline{y}) \underline{P}(\underline{y}) d\underline{y} \|^2 \} \\
 &+ \text{Area}(A) \int_A \| (-\epsilon) \underline{M}(\underline{x}, \underline{y}) \|^2 d\underline{y} \\
 &= E \{ \| \underline{P}(\underline{x}) - \int_A \underline{H}^{(0)}(\underline{x}, \underline{y}) \underline{r}(\underline{y}) d\underline{y} \|^2 \} - \epsilon \int_A \| \underline{M}(\underline{x}, \underline{z}) \|^2 d\underline{z}
 \end{aligned}$$

$$+ \epsilon^2 \{ E \{ \left\| \int_A \underline{M}(\underline{x}, \underline{y}) \underline{P}(\underline{y}) d\underline{y} \right\|^2 \} + \text{Area}(A) \int_A \left\| \underline{M}(\underline{x}, \underline{y}) \right\|^2 d\underline{y} \} \quad (27)$$

Assuming that  $\underline{H}^{(0)}(\underline{x}, \underline{y})$  is not a solution of the Wiener-Hopf equation and remembering the definition of  $\underline{M}(\underline{x}, \underline{y})$ , (25) makes it clear that

$$- \epsilon \int_A \left\| \underline{M}(\underline{x}, \underline{z}) \right\|^2 d\underline{z} < 0$$

Thus, for  $\epsilon$  sufficiently small (27) shows that

$$E \{ \left\| \underline{P}(\underline{x}) - \int_A \underline{H}(\underline{x}, \underline{y}) \underline{r}(\underline{y}) d\underline{y} \right\|^2 \} < E \{ \left\| \underline{P}(\underline{x}) - \int_A \underline{H}^{(0)}(\underline{x}, \underline{y}) \underline{r}(\underline{y}) d\underline{y} \right\|^2 \}$$

where, by (24) and (25),

$$\begin{aligned} \underline{H}(\underline{x}, \underline{z}) &= \underline{H}^{(0)}(\underline{x}, \underline{z}) + \underline{\Delta}(\underline{x}, \underline{z}) \\ &= \underline{H}^{(0)}(\underline{x}, \underline{z}) - \epsilon E \left[ \left( \underline{P}(\underline{x}) - \int_A \underline{H}^{(0)}(\underline{x}, \underline{y}) \underline{r}(\underline{y}) d\underline{y} \right) \underline{r}^t(\underline{z}) \right] \end{aligned}$$

Hence, it has been shown that if  $\underline{H}^{(0)}(\underline{x}, \underline{y})$  is not a solution to the Wiener-Hopf equation  $\underline{H}^{(0)}(\underline{x}, \underline{y})$  does not minimize the mean-square error.

#### 4.0 SOLUTION UNDER THE ASSUMPTION OF STATIONARITY AND NON-CAUSALITY

If we allow the matrix filter to be non-causal and further assume spatial stationarity, we find the result,

$$\tilde{H}(\omega_x, \omega_y) = \tilde{T}^t \tilde{G}(\omega_x, \omega_y) \tilde{T} \quad (28)$$

where

$$\tilde{G}(\omega_x, \omega_y) = \begin{bmatrix} \frac{\lambda_1 S(\omega_x, \omega_y)}{\lambda_1 S(\omega_x, \omega_y) + \sigma_n^2} & & \\ & \ddots & \\ & & \frac{\lambda_N S(\omega_x, \omega_y)}{\lambda_N S(\omega_x, \omega_y) + \sigma_N^2} \end{bmatrix} \quad (29)$$

and  $S(\omega_x, \omega_y)$  is the spatial power spectral density of the original bands, and  $\sigma_N^2$  is the noise variance. This matrix filter consists of  $N^2$  filters given by

$$H_{i,j}(\omega_x, \omega_y) = \sum_{k=1}^N e_{k,i} e_{k,j} \left( \frac{\lambda_n S(\omega_x, \omega_y)}{\lambda_n S(\omega_x, \omega_y) + \sigma_n^2} \right) \quad (30)$$

where  $e_{k,i}$  is the  $k,i$ -th element of  $\tilde{T}$  ( $i$ -th element of the  $k$ -th eigenvector of  $\tilde{C}$ ).

Implementation of the  $N^2$  filters requires

$$O_H = N^2 N_S^2 (1 + 4 \log_2 N_S)$$

operations (multiplications and additions), where  $N_S$  is the number of samples per line in each of the original (square) images. If we implement the filters using the  $\tilde{G}$  matrix preceded and proceeded by the K-L transform as indicated in (28), then the number of operations is

$$O_{K-L} = N N_S^2 [2N + (1 + 4 \log_2 N_S)] .$$

Clearly,

$$\frac{O_{K-L}}{O_H} < 1 \text{ for } N \geq 2, N_S \geq 2.$$

In fact,

$$\lim_{N_S \rightarrow \infty} \frac{O_{K-L}}{O_H} = \frac{1}{N}$$

and the filter implementation in (28) results in an approximate  $1/N$  reduction in the number of operations for typical pictures ( $N_S \geq 256$ ). For example, with  $N_S = 256$  and  $N = 3$  (color data),  $O_G/O_H = 1/2.54$  ( $7.68 \times 10^6$  ops versus  $19.5 \times 10^6$  ops).

#### 5.0 CLASS-DEPENDENT K-L TRANSFORMATION

The presence of linear class structure within multiband data suggests the use of K-L transforms matched to the covariance matrices of each class. In so doing we achieve a piece wise linear approximation to a non-linear transformation of the multiband data, and approximate the "intrinsic dimensionality" of the multiband data. The result is a further compression of data variance into a smaller number of principal components compared to the global K-L based on a single covariance matrix computed over the entire picture.

Our approach to classification is based on a simple unsupervised clustering algorithm in which an initial (random) guess of both the number of classes and their mean vectors are iterated until the mean vectors remain constant and the inter class distances (Bhattacharyya distance) satisfy a threshold. Upon convergence of the algorithm we have the number of class and their covariance matrices. This information is used to compute the required set of K-L transforms, and to implement a conventional likelihood-ratio hypothesis test for each data vector comprising the multiband image.

Once the data is classified we then perform the class dependent K-L transformation. The MSE-optimum spatial filters are applied



to the principal component images, followed by inverse transformation.

The effectiveness of the above procedure has been verified using a color test image (3 bands: red, green, and blue) of an F-16 aircraft obtained from the USC Image Processing Institute.

In Figure 2 we show histograms (estimates of the probability density function) of a) the original data, b) global principal components and c) the class dependent principal components. Note the significant reduction in variance in c).

For this particular data set the clustering algorithm generated five classes. These classes are depicted in Figure 3. The uniqueness of each class is somewhat evident in Figures (4) and (5) where we show a two dimensional projection of the three dimensional joint density function for the original bands. Note the varying directions of maximum variance for each class, and therefore, the appropriateness of separate K-L transform for each class.




Fig. 2--Comparison of density functions




Fig. 3--Map of estimated classes




Fig. 4--Contour plot of red-green joint density




Fig. 5--Three dimensional view of Fig. (4)